

EDITORIAL

Hepatocentric Leptin Signaling Modulates Gluconeogenesis via MKP-3

Since its discovery in 1994, leptin has been recognized as a satiety hormone required for body weight homeostasis. Leptin is secreted predominantly by white adipose tissue, and its levels in blood are correlated positively with the amount of body fat. Extensive studies of leptin's actions in the central nervous system (CNS) have shown its ability to control food intake and energy expenditure. However, despite the profound obesity and diabetes resulting from homozygous loss of leptin or its receptor, there has been very limited efficacy of leptin treatment for obesity because most obese individuals already have high circulating leptin levels, rendering them unresponsive to its weight-reducing effects.

Research now has shifted to leptin's effects in the periphery, particularly in the context of its glucoregulatory actions, where it may have a role independent of body weight regulation.¹ Indeed, hyperinsulinemia occurs before weight gain in leptin-deficient *ob/ob* mice,² and there are significant improvements in hyperglycemia and hyperinsulinemia before weight loss in leptin-treated *ob/ob* mice.³

The leptin receptor (Obr) is present at highest levels in the CNS, but also is expressed throughout the periphery. There are 6 isoforms of Obr that result from alternative splicing and, importantly, only *Obrb*, the long leptin receptor isoform, is capable of mediating signal transduction. *OBRb* activates the JAK-STAT3 and PI3K systems, which are critical pathways involved in energy homeostasis and glucose metabolism, respectively.

In this issue of *Cellular and Molecular Gastroenterology and Hepatology*, Huang et al and He et al reported hepatocyte-specific effects of leptin signaling through *Obrb* to suppress glucose production. The investigators showed that leptin treatment of primary hepatocytes and hepatoma cells in vitro resulted in STAT3 phosphorylation, suppression of glucose production, and decreased expression of the gluconeogenic genes *G6pc*, *Pepck1*, and *Pgc1a*. These effects were reversed after small interfering RNA-mediated suppression of *Obr*. These in vitro findings support a cell-autonomous effect of leptin on hepatocyte glucose production (Figure 1). In 2 different mouse models of obesity—leptin-receptor-deficient *db/db* mice and high-fat diet-fed mice—*Obrb* overexpression specifically in liver was sufficient to lower blood glucose levels, improve glucose tolerance, and improve insulin tolerance. On the other hand, small interfering RNA-mediated suppression of liver *Obr* in lean mice had no effect on blood glucose.

Through which signaling pathway does hepatocyte leptin signaling control glucose production? Previously published

data from the same research group showed that mitogen-activated protein kinase phosphatase-3 (MKP-3) is increased significantly in the liver of diet-induced obese mice and has regulatory control over gluconeogenesis.⁴ They showed that MKP-3 dephosphorylates FoxO1 to promote its nuclear translocation, subsequently inducing the transcription of gluconeogenic genes.⁴ The investigators now show that leptin and *Obrb* overexpression in the presence of leptin significantly decreases MKP-3 protein levels in primary hepatocytes and in mice, whereas *Obrb* suppression in primary hepatocytes increases MKP-3. Moreover, *Mkp-3* deficiency blocks the ability of leptin and *Obrb* overexpression to suppress glucose production and gluconeogenic gene expression, showing that MKP-3 mediates the effects of leptin signaling on hepatic gluconeogenesis (Figure 1).

These data support an effect of leptin signaling through STAT3 and MKP-3 to decrease gluconeogenesis in




Figure 1. Model of hepatic leptin suppression of gluconeogenesis. Leptin binds to its receptor *OBRb* at the plasma membrane, which activates STAT3 via JAK2 signaling. STAT3, possibly by complexing directly with ERK1/2 and MKP-3, leads to MKP-3 protein degradation. Decreased MKP-3 levels lead to increased phosphorylation of FOXO1, thereby excluding FOXO1 from the nucleus and decreasing expression of key gluconeogenic genes, *G6pc* and *Pck1*, and their regulatory gene *Pgc1a*. Created with BioRender.com

117 hepatocytes. However, critical questions remain. The in-
 118 vestigators noted no effect of liver *Obr* suppression in lean
 119 mice. Is hepatocyte leptin signaling activated only under
 120 certain physiologic or pathologic conditions? Leptin can be
 121 thought of as an adipostat, relaying information about body
 122 fat status to the brain to control energy balance. Is there a
 123 role for leptin's effects on hepatic glucose production in this
 124 context? How do the effects of leptin compare, or synergize,
 125 with classic suppression of gluconeogenesis by insulin
 126 signaling? Overall, this work highlights the signaling role of
 127 leptin outside the CNS and suggests that further research
 128 should be performed to understand these pathways.

129
 130 *JENNIFER R. GAMARRA, MD*

131 *REBECCA A. HAEUSLER, MD*

132 Naomi Berrie Diabetes Center, Department of Pathology and
 133 Cell Biology
 134 Columbia University Medical Center
 135 New York, New York

137 References

- 138 1. D'souza AM, Neumann UH, Glavas MM, Kieffer TJ. The
 139 glucoregulatory actions of leptin. *Mol Metab* 2017;
 140 6:1052–1065.
- 141 2. Genuth SM, Przybylski RJ, Rosenberg DM. Insulin
 142 resistance in genetically obese, hyperglycemic mice.
 143 *Endocrinology* 1971;88:1230–1238.
 144
 145
 146
 147
 148
 149
 150
 151
 152
 153
 154
 155

3. Harris RBS, Zhou J, Redmann SM, Smagin GN, 156
 Smith SR, Rodgers E, Zachwieja JJ. A leptin dose- 157
 response study in obese (ob/ob) and lean (+/?) mice. 158
Endocrinology 1998;139:8–19. 159
4. Wu Z, Jiao P, Huang X, Feng B, Feng Y, Yang S, 160
 Hwang P, Du J, Nie Y, Xiao G, Xu H. MAPK 161
 phosphatase-3 promotes hepatic gluconeogenesis 162
 through dephosphorylation of forkhead box O1 in mice. 163
J Clin Invest 2010;120:3901–3911. 164

165 Correspondence

166 Address correspondence to: Rebecca A. Haeusler, MD, Naomi Berrie Diabetes
 167 Center, Department of Pathology and Cell Biology, Columbia University
 168 Medical Center, 1150 Saint Nicholas Avenue, Room 303A, New York, New
 169 York 10032. e-mail: rah2130@cumc.columbia.edu. Q23 170

171 Conflicts of interest

172 The authors disclose no conflicts. Q4 173

174 Funding

175 Supported by a NSF-GRFP fellowship (J.R.G.); National Institutes of Health
 176 grants R01DK115825 and R01HL125649, American Diabetes Association
 177 grant 7-20-IBS-130 (R.A.H.), and National Institute of Diabetes and Digestive
 178 and Kidney Diseases Digestive Disease Research Center support grant P30
 179 DK132710. Q6 175 Q7 178

180 © 2022 The Authors. Published by Elsevier Inc. on behalf of the AGA
 181 Institute. This is an open access article under the CC BY-NC-ND license
 182 (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).
 183 2352-345X
 184 <https://doi.org/10.1016/j.jcmgh.2022.08.006>
 185
 186
 187
 188
 189
 190
 191
 192
 193
 194